#### ACETAMINOPHEN DOSAGE CHART

The proper dosage is determined by weight not age. If your child is not old enough to stand on a scale you can weigh yourself and then weigh holding your child and subtract the two.

Child's Weight	6 to 11 lbs	12 to 17 lbs	18 to 23 lbs	24 to 35 lbs	36 to 47 lbs	48 to 59 lbs	60 to 71 lbs	72 to 95 lbs	96+ lbs
Infant Drops	1.25ml in syringe	2.5ml in syringe	3.75ml in syringe	5ml in syringe					
Children's Liquid		<sup>1</sup> / <sub>2</sub> tsp (2.5ml in cup)	<sup>3</sup> / <sub>4</sub> tsp (3.75ml in cup)	1 tsp (5ml in cup)	1 <sup>1</sup> / <sub>2</sub> tsp (7.5ml in cup)	2 tsp (10ml in cup)	$2^{1}/_{2}$ tsp (12.5ml on cup)	3 tsp (15ml in cup)	4 tsp (20ml in cup)
Children's Chew/Meltaways			11/2	2	3	4	21/2	6	8
Junior Strength Chew/Meltaways				1	11/2	2	21/2	3	4
Adult Regular- Strength Tabs (325mg)							1	1 to 1 <sup>1</sup> / <sub>2</sub>	2

## You may repeat the dose every 4 hours while symptoms last

#### **Active Ingredients**

Infant Drops: 160mg/5ml per included syringe (shake well before

using)

Children's Liquid: 160mg/5ml per teaspoon (shake well before using)

Children's Chew/Meltaways: 80mg each Junior Strength Chew/Meltaways: 160mg each

Adult Regular Strength Tablets: 325mg

## **Safety Tips for Acetaminophen**

- Don't give to a baby under 3 months without a doctors approval
- Always use the measuring device that comes with the medication
- Never give it to a child who is taking another medication that contains acetaminophen without a doctors approval
- Do not exceed 5 doses in 24 hours

#### IBUPROFEN DOSAGE CHART

The proper dosage is determined by weight not age. If your child is not old enough to stand on a scale you can weigh yourself and then weigh holding your child and subtract the two.

Child's Weight	6 to 11 lbs	12 to 17 lbs	18 to 23 lbs	24 to 35 lbs	36 to 47 lbs	48 to 59 lbs	60 to 71 lbs	72 to 95 lbs	96+ lbs
Children's Liquid		¹/₂tsp	³/ <sub>4</sub> tsp	1 tsp	1 <sup>1</sup> / <sub>2</sub> tsp	2 tsp	2 <sup>1</sup> / <sub>2</sub> tsp	3 tsp	4 tsp
Children's Chew					3	4	5	6	8
Junior Strength Chew/Caplets						2	<b>2</b> <sup>1</sup> / <sub>2</sub>	3	4
Adult Regular Strength Tabs							1	1 to 1 <sup>1</sup> / <sub>2</sub>	2

# You may repeat the dose every 6-8 hours while symptoms last

### **Active Ingredients**

Children's Liquid: 100mg (5ml) per teaspoon (shake well before

using)

Children's Chewables: 50mg each

Junior Strength Chew/Caplets: 100mg each Adult Regular Strength Tablets: 200mg each

## **Safety Tips for Ibuprofen**

- Don't give to a child under 6 months of age without a doctor's approval
- Always use the measuring device that comes with the medication
- Never give it to a child who is taking another medication that contain NSAIDs without a doctors approval
- Do not exceed 4 doses in 24 hours